



www.yourgicenter.com
 facebook.com/yourgicenter
 Email: hello@yourgicenter.com
 Toll free: 888-292-0010
 Fax: 979-292-0488

3 DAY PREPOPIK PREPARATION FOR COLONOSCOPY

Patient Check-In Time _____ Patient Procedure Date _____

(Please note: Patient Check-In Time is not the procedure time! It is merely the time we wish for you to begin the check in process that starts with signing in at the front window.)

Location where your procedure will take place:

- The G.I.Center, 109 Parking Way, Lake Jackson, TX 77566**
 *The front door opens at 6:30 a.m.
 *If you need to cancel or reschedule your procedure, please call 979-292-0033 or 979-285-9948
 *Our office requires a 24 hours notice if you are going to cancel or reschedule your appointment otherwise a \$100.00 cancellation fee will be billed to you.

- Your GI Center at Pearland (Pearland Premier Surgery Center)**
2813 Smith Ranch Road, Pearland, TX 77584
 * The front door opens at 7:00 a.m.
 *If you need to cancel or reschedule your procedure, please call (713) 436-8171
 *Our office requires a 24 hours notice if you are going to cancel or reschedule your appointment otherwise a \$100.00 cancellation fee will be billed to you.

- Hospital * 24 hour notice still required to avoid \$100.00 fee for rescheduled or cancelled procedures.**

Patient Checklist:

- Purchase four Dulcolax laxative tablets (over the counter) and Prepopik (prescription given) from your local pharmacy.
- You MUST have a ride to and from the facility** because you will not be allowed to drive until 12 hours after the procedure. This is because anesthesia will be given to you to sedate you during your procedure. **If your ride must leave the facility, we must have a number (such as a cell phone number) to reach them on. If you do not have a ride in a timely manner, a cab may be called for you at your expense at the discretion of our ASC Nursing Staff.**
- Prescription blood thinners must be stopped prior to the procedure. (Please see the chart below)** (*Aspirin is okay to take unless specifically instructed otherwise*). All other medications can be taken on its normal schedule with a small sip of water.

5 days	2 days
Coumadin, Warfarin, Plavix	Xarelto, Aggrenox, Eliquis, Pradaxa
<i>*Aspirin is okay to take unless specifically instructed otherwise.*</i>	

- DIABETIC PATIENTS:** No Diabetic medication including pills and injections on the day of the procedure
- if you use a Nebulizer,** please do a breathing treatment before coming for your procedure.
- If you use an inhaler,** bring it with you the day of your procedure.

www.yourgicenter.com

Pearland
 2813 Smith Ranch Rd
 Pearland, TX 77584
 713-436-8171

Friendswood
 1305 Parkwood Ave
 Friendswood, TX 77546
 832-808-9872

Lake Jackson
 109 Parking Way
 Lake Jackson, TX 77566
 979-292-0033

Bay City
 720 Avenue F North
 Bay City, TX 77414
 979-292-0033



www.yourgicenter.com
facebook.com/yourgicenter
Email: hello@yourgicenter.com
Toll free: 888-292-0010
Fax: 979-292-0488

INSTRUCTIONS ON YOUR PREPARATION FOR THE COLONOSCOPY.

- **Three days before Colonoscopy:**
 - Take 2 Dulcolax tabs by mouth by 5 PM
 - Regular Diet this day.
- **Two days before Colonoscopy:**
 - Take 2 Dulcolax tabs by mouth by 5 PM
 - Start Clear Liquid Diet this day (no solid foods or milk products).
- **The day before your Colonoscopy:**
 - Continue with a Clear Liquid Diet
 - At 7 PM drink one dose of Prepopik followed by five, 8 ounce glasses of clear liquid.
Make sure to mix Prepopik for 4 minutes continuously.
- **The day of your Colonoscopy:**
 - Take the 2nd dose of Prepopik followed by three glasses of water.
Once again being sure to mix Prepopik for 4 minutes continuously.
Complete this AT LEAST 4 hours before your check in time, with nothing further by mouth (no liquids, mints, gum, chewing tobacco, etc.).

*Clear liquids include all of the following that are not colored red or purple. Strained fruitjuices **without** pulp only (apple, white grape, lemonade), water, clear broth or bouillon, coffee (without milk or nondairy creamer), Gatorade, carbonated and noncarbonated soft drinks. Kool-Aid, (or other fruit flavored drinks), plain jello (without added fruit or topping) and ice popsicles. Look under tab for DIETS in our website at www.yourgicenter.com and get detail list of foods for Clear Liquid Diets

**Bowel movements should be liquid and without particles/pieces of solid stool.

- **SEE ATTACHED INFORMMATION FOR CLEAR LIQUID DIET.**
- **IF YOU ARE EXPERIENCING LOW SUGAR SYMPTOMS, IMMEDIATELY TAKE SOME SUGAR SOLUTION OR CLEAR JUICE SUCH AS APPLE JUICE.**
- **NO SMOKING AFTER MIDNIGHT.**
- **DO NOT TAKE FISH OIL, VITAMIN A, OR ANY OTHER OIL BASED MEDICATION OR VITAMINS FOR 3 DAYS PRIOR TO YOUR PROCEDURE.**

www.yourgicenter.com

Pearland
2813 Smith Ranch Rd
Pearland, TX 77584
713-436-8171

Friendswood
1305 Parkwood Ave
Friendswood, TX 77546
832-808-9872

Lake Jackson
109 Parking Way
Lake Jackson, TX 77566
979-292-0033

Bay City
720 Avenue F North
Bay City, TX 77414
979-292-0033



www.yourgicenter.com
 facebook.com/yourgicenter
 Email: hello@yourgicenter.com
 Toll free: 888-292-0010
 Fax: 979-292-0488

CLEAR LIQUID DIET

This diet provides fluids that leave little residue and are easily absorbed with minimal digestive activity. This diet is inadequate in all essential nutrients and is recommended only if clear liquids are temporarily needed. No red or purple liquids should be consumed!

Food Group	Foods Allowed	Foods to Avoid
Milk & beverages <i>No red or purple liquids!</i>	Tea (decaffeinated or regular), carbonated beverages, fruit flavored drinks	Milk, Milk drinks
Meats & meat substitutes	None	All
Vegetables	None	All
Fruits & fruit juices	Strained fruit juices: apple, white grape, lemonade	Fruit juices with unstrained fruit
Grains & starches	None	All
Soups	Clear broth, consommé	All others
Desserts	Clear flavored gelatin, popsicles (no red or purple flavors)	All others
Fats	None	All
Miscellaneous	Sugar, honey, syrup, clear hard candy, salt	All others
Examples of Clear Liquid Diet:		
Breakfast	Lunch	Dinner
White grape juice	Apple Juice	Lemonade
Clear broth	Clear broth	Clear broth
Jell-O ^{®*}	Jell-O ^{®*}	Jell-O ^{®*}
Tea	Tea	Tea
<i>*plain only, no fruit or toppings</i>	<small>Jell-O is a registered trademark of Kraft General Foods, Inc.</small>	

www.yourgicenter.com

Pearland
 2813 Smith Ranch Rd
 Pearland, TX 77584
 713-436-8171

Friendswood
 1305 Parkwood Ave
 Friendswood, TX 77546
 832-808-9872

Lake Jackson
 109 Parking Way
 Lake Jackson, TX 77566
 979-292-0033

Bay City
 720 Avenue F North
 Bay City, TX 77414
 979-292-0033



www.yourgicenter.com
facebook.com/yourgicenter
Email: hello@yourgicenter.com
Toll free: 888-292-0010
Fax: 979-292-0488

Insulin Patients

1. For Short/ultra short acting insulin – skip morning dose
2. For long acting – Take half dose (in the morning or the night before the procedure)
3. For Combination / Intermediate – Take half dose in the morning
4. Skip all morning diabetic pills until after the procedure.

w w w . y o u r g i c e n t e r . c o m

Pearland
2813 Smith Ranch Rd
Pearland, TX 77584
713-436-8171

Friendswood
1305 Parkwood Ave
Friendswood, TX 77546
832-808-9872

Lake Jackson
109 Parking Way
Lake Jackson, TX 77566
979-292-0033

Bay City
720 Avenue F North
Bay City, TX 77414
979-292-0033